

MANE 3332.04

Section 1

MANE 3332.04

## Subsection 1

### Chapter 4 Content

## Continuous Random Variable

- The **probability distribution** of a random variable  $X$  is a description of the set of probabilities associated with the possible values of  $X$
- Density functions are commonly used in engineering to describe physical systems.
- A **probability density function**  $f(x)$  can be used to describe the probability distribution of a continuous random variable

*pdf versus pmf (chapter 3)*

### Probability Density Function

- Notice the difference from a discrete random variable
- The formal definition of a probability density function is a function such that
  - 1  $f(x) \geq 0$
  - 2  $\int_{-\infty}^{\infty} f(x) dx = 1$
  - 3  $P(a \leq X \leq b) = \int_a^b f(x) dx$

*Handwritten notes:*


Pmf  
~~1~~  $f(x) \geq 0$   
②  $\sum_{\text{all } x} f(x) = 1$

### Probability Density Function

- Any interesting property of continuous random variables is

$$P(x_1 \leq X \leq x_2) = P(x_1 < X \leq x_2) = P(x_1 \leq X < x_2) = P(x_1 < X < x_2)$$

- Does not apply to discrete random variables
- Explanation



graph on the right does not include the exact value of  $x_1$

### Cumulative Distribution Function

The cumulative distribution function for a continuous random variable  $X$  is

$$F(x) = P(X \leq x) = \int_{-\infty}^x f(y) dy$$

### Mean and Variance of a Continuous Random Variable

- The mean value of a continuous random variable is defined to be

$$\mu = E(X) = \int_{-\infty}^{\infty} xf(x) dx$$

$$\mu = \sum_{\text{all } y} xf(y)$$

- The variance of a continuous random variable is defined to be

$$\begin{aligned} \sigma^2 &= V(X) = E(X - \mu)^2 \\ &= \int_{-\infty}^{\infty} (x - \mu)^2 f(x) dx = \int_{-\infty}^{\infty} x^2 f(x) dx - \mu^2 \end{aligned}$$

- The standard deviation of  $X$  is


$$\sigma = \sqrt{V(X)}$$

## Continuous Uniform Distribution

The continuous uniform distribution is the analog of the discrete uniform distribution in that all outcomes are equally likely to occur

- A continuous uniform distribution for the random variable  $X$  has a probability density function

$$f(x) = \frac{1}{b-a}, \quad a \leq x \leq b$$



- The mean of the uniform distribution is

$$\mu = E(X) = \frac{a+b}{2}$$

Handwritten derivation for the mean:

$$\int_{-\infty}^{\infty} x f(x) dx = \int_{-\infty}^{\infty} x \left( \frac{1}{b-a} \right) dx$$

$$= \frac{1}{b-a} \left[ \int_{-\infty}^a 0 dx + \int_a^b x dx + \int_b^{\infty} 0 dx \right]$$

- The variance of  $X$  is

$$\sigma^2 = V(X) = \frac{(b-a)^2}{12}$$

Handwritten derivation for the variance:


$$= \frac{1}{b-a} \int_a^b x^2 dx$$

$$= \left( \frac{1}{b-a} \right) \frac{x^3}{3} \Big|_a^b$$

$$= \left( \frac{1}{b-a} \right) \left( \frac{b^3}{3} - \frac{a^3}{3} \right)$$

$$= \frac{1}{2} \left( \frac{1}{b-a} \right) (a+b)(a-b)$$






$$f(x) = 2 \quad \text{for } 49.75 < x < 50.25$$

$$P(x > 50) = \int_{50}^{50.25} 2 \, dx = 2x \Big|_{x=50}^{50.25}$$

Uniform Problem 4.1.6

- See page P-25

$$= 2(50.25 - 50) = .5$$



$$P(x < 49.8) = \int_{49.75}^{49.8} 2 \, dx = 2x \Big|_{x=49.75}^{49.8} = 2(49.8 - 49.75) = .1$$

$$P(x < 49.8) = F(49.8)$$

$$P(x > 50) = 1 - F(50)$$

9 / 42

e

Attendance 1-D

# Gaussian (Gauss)

MANE 3332.04 Chapter 4 Content

## The Normal Distribution

- The normal distribution is the most widely used and important distribution in statistics.
- You must master this!
- A random variable  $X$  with probability density function




$$f(x) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{(x-\mu)^2}{2\sigma^2}} \text{ for } -\infty < x < \infty$$

has a **normal distribution** with parameters  $\mu$  and  $\sigma$  where  $-\infty < \mu < \infty$  and  $\sigma > 0$

- The normal distribution with parameters  $\mu$  and  $\sigma$  is denoted  $N(\mu, \sigma^2)$
- An interesting web-site is <http://www.seeingstatistics.com/seeingTour/normal/shape3.html>

10 / 42

give  $N =$   
 $\sigma_1 =$   
 $\sigma_2 =$   
 $\sigma_3 =$



MANE 3332.04 Chapter 4 Content

## Mean and Variance of the Normal Distribution

- The mean of the normal distribution with parameters  $\mu$  and  $\sigma$  is

$$E(X) = \mu$$

- The variance of the normal distribution with parameters  $\mu$  and  $\sigma$  is

MANE 3332.04 Chapter 4 Content

- The variance of the normal distribution with parameters  $\mu$  and  $\sigma$  is

$$V(X) = \sigma^2$$

11 / 42

MANE 3332.04 Chapter 4 Content

### Central Limit Theorem

- Brief introduction
- States that the distribution of the average of independent random variables will tend towards a normal distribution as  $n$  gets large
- More details later

*n around 25-30*

12 / 42

## Calculating Normal Probabilities

- Is somewhat complicated
- The difficulty is  $\int_a^b f(x) dx$  does not have a closed form solution
- Probabilities must be found by numerical techniques (tabled values)
- It is very helpful to draw a sketch of the desired probabilities (I require this)

How to use without closed form  
Solution

- ① Tables
- ② Computers

closed form  
recall uniform

$$\int_a^x f(u) du = \frac{x-a}{b-a}$$


answer does not contain  
an integral

## The Standard Normal Distribution

- A normal random variable with  $\mu = 0$  and  $\sigma = 1$  is called a **standard normal** random variable
- A standard normal random variable is denoted as  $z$
- The cumulative distribution function for a standard normal is defined to be the function  $F(x)$  for standard normal is  $\Phi(x)$

$$\Phi(z) = P(Z \leq z) = F(z)$$

- These probabilities are contained in Appendix Table III on pages A-8 and A-9



$\Phi$  - capital phi

## Cumulative Standard Normal Distribution

A.8 APPENDIX A Statistical Tables and Charts




TABLE II Cumulative Standard Normal Distribution

z	-0.09	-0.08	-0.07	-0.06	-0.05	-0.04	-0.03	-0.02	-0.01	0.00
-3.9	0.00003	0.00004	0.00005	0.00007	0.00009	0.00011	0.00014	0.00018	0.00023	0.00029
-3.8	0.00006	0.00008	0.00010	0.00013	0.00016	0.00020	0.00025	0.00031	0.00038	0.00046
-3.7	0.00011	0.00014	0.00017	0.00021	0.00026	0.00032	0.00039	0.00047	0.00056	0.00066
-3.6	0.00017	0.00021	0.00026	0.00031	0.00037	0.00044	0.00052	0.00061	0.00071	0.00082
-3.5	0.00024	0.00029	0.00035	0.00041	0.00048	0.00056	0.00064	0.00073	0.00083	0.00094
-3.4	0.00032	0.00038	0.00045	0.00052	0.00060	0.00068	0.00077	0.00086	0.00096	0.00106
-3.3	0.00040	0.00047	0.00055	0.00063	0.00071	0.00080	0.00089	0.00098	0.00108	0.00118
-3.2	0.00049	0.00057	0.00065	0.00073	0.00082	0.00091	0.00100	0.00109	0.00119	0.00129
-3.1	0.00058	0.00067	0.00075	0.00084	0.00093	0.00102	0.00111	0.00120	0.00129	0.00139
-3.0	0.00068	0.00077	0.00085	0.00094	0.00103	0.00112	0.00121	0.00130	0.00139	0.00148
-2.9	0.00078	0.00087	0.00096	0.00104	0.00113	0.00122	0.00131	0.00140	0.00148	0.00157
-2.8	0.00088	0.00097	0.00106	0.00114	0.00123	0.00132	0.00141	0.00149	0.00157	0.00166
-2.7	0.00098	0.00107	0.00116	0.00125	0.00134	0.00143	0.00151	0.00160	0.00168	0.00177
-2.6	0.00108	0.00117	0.00126	0.00135	0.00144	0.00152	0.00161	0.00169	0.00178	0.00187
-2.5	0.00118	0.00127	0.00136	0.00145	0.00154	0.00162	0.00171	0.00179	0.00188	0.00196
-2.4	0.00128	0.00137	0.00146	0.00155	0.00164	0.00173	0.00181	0.00190	0.00198	0.00206
-2.3	0.00138	0.00147	0.00156	0.00165	0.00174	0.00182	0.00191	0.00199	0.00208	0.00216
-2.2	0.00148	0.00157	0.00166	0.00175	0.00184	0.00192	0.00201	0.00209	0.00218	0.00226
-2.1	0.00158	0.00167	0.00176	0.00185	0.00193	0.00202	0.00210	0.00219	0.00227	0.00235
-2.0	0.00168	0.00177	0.00186	0.00194	0.00203	0.00211	0.00220	0.00228	0.00236	0.00244
-1.9	0.00178	0.00187	0.00195	0.00204	0.00212	0.00221	0.00229	0.00237	0.00245	0.00253
-1.8	0.00188	0.00196	0.00205	0.00213	0.00221	0.00230	0.00238	0.00246	0.00254	0.00262
-1.7	0.00197	0.00206	0.00214	0.00222	0.00230	0.00238	0.00246	0.00254	0.00262	0.00270
-1.6	0.00207	0.00215	0.00223	0.00231	0.00239	0.00247	0.00255	0.00263	0.00271	0.00279
-1.5	0.00217	0.00225	0.00232	0.00240	0.00248	0.00256	0.00264	0.00271	0.00279	0.00287
-1.4	0.00227	0.00234	0.00242	0.00250	0.00258	0.00266	0.00273	0.00281	0.00289	0.00296
-1.3	0.00237	0.00244	0.00252	0.00260	0.00268	0.00275	0.00283	0.00290	0.00298	0.00305
-1.2	0.00247	0.00254	0.00262	0.00270	0.00277	0.00285	0.00292	0.00299	0.00306	0.00313
-1.1	0.00257	0.00264	0.00271	0.00279	0.00286	0.00293	0.00300	0.00307	0.00314	0.00321
-1.0	0.00267	0.00274	0.00281	0.00288	0.00295	0.00302	0.00309	0.00315	0.00322	0.00328
-0.9	0.00276	0.00283	0.00290	0.00297	0.00304	0.00311	0.00317	0.00324	0.00330	0.00336
-0.8	0.00286	0.00292	0.00299	0.00306	0.00312	0.00319	0.00325	0.00332	0.00338	0.00344
-0.7	0.00296	0.00302	0.00309	0.00315	0.00321	0.00328	0.00334	0.00340	0.00346	0.00352
-0.6	0.00306	0.00312	0.00318	0.00324	0.00330	0.00336	0.00342	0.00348	0.00354	0.00359
-0.5	0.00315	0.00321	0.00327	0.00333	0.00339	0.00345	0.00351	0.00356	0.00362	0.00367
-0.4	0.00325	0.00331	0.00337	0.00343	0.00349	0.00354	0.00360	0.00365	0.00371	0.00376
-0.3	0.00334	0.00340	0.00346	0.00352	0.00357	0.00363	0.00368	0.00373	0.00379	0.00384
-0.2	0.00344	0.00349	0.00355	0.00360	0.00365	0.00370	0.00375	0.00380	0.00385	0.00390
-0.1	0.00354	0.00359	0.00364	0.00369	0.00374	0.00379	0.00384	0.00389	0.00394	0.00398
0.0	0.00359	0.00364	0.00369	0.00374	0.00379	0.00384	0.00389	0.00394	0.00398	0.00403

### Cumulative Standard Normal Distribution

APPENDIX A Statistical Tables and Charts A-9

$$\Phi(z) = P\{Z \leq z\} = \int_{-\infty}^z \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}a^2} da$$



**TABLE III** Cumulative Standard Normal Distribution (continued)

	0.0	0.1	0.2	0.3	0.4	0.5	0.6	0.7	0.8	0.9
10	5.50000	5.50498	5.50976	5.51467	5.51953	5.52445	5.52932	5.53420	5.53911	5.54406
11	5.53898	5.54387	5.54878	5.55371	5.55870	5.56368	5.56869	5.57400	5.57842	5.58343
12	5.57920	5.58410	5.58904	5.59400	5.59898	5.60398	5.60899	5.61402	5.61908	5.62415
13	5.62423	5.62928	5.63436	5.63946	5.64458	5.64971	5.65486	5.65993	5.66502	5.67013
14	5.65452	5.65957	5.66467	5.66980	5.67494	5.67991	5.68493	5.68997	5.69498	5.69999
15	5.69442	5.69947	5.70456	5.70966	5.71478	5.71991	5.72505	5.73019	5.73534	5.74049
16	5.73498	5.74004	5.74512	5.75021	5.75531	5.76042	5.76554	5.77067	5.77580	5.78094
17	5.77508	5.78014	5.78523	5.79033	5.79544	5.79999	5.80505	5.80999	5.81498	5.81999
18	5.78145	5.78650	5.79157	5.79667	5.80178	5.80689	5.81199	5.81709	5.82219	5.82729
19	5.81633	5.82143	5.82654	5.83165	5.83676	5.84187	5.84697	5.85208	5.85718	5.86229
20	5.84145	5.84655	5.85166	5.85676	5.86187	5.86697	5.87208	5.87718	5.88229	5.88739
21	5.87646	5.88156	5.88667	5.89177	5.89687	5.90197	5.90708	5.91218	5.91729	5.92239
22	5.88940	5.89450	5.89960	5.90470	5.90980	5.91490	5.92000	5.92510	5.93020	5.93530
23	5.90309	5.90819	5.91329	5.91839	5.92349	5.92859	5.93369	5.93879	5.94389	5.94899
24	5.93189	5.93699	5.94209	5.94719	5.95229	5.95739	5.96249	5.96759	5.97269	5.97779
25	5.93193	5.93703	5.94213	5.94723	5.95233	5.95743	5.96253	5.96763	5.97273	5.97783
26	5.94205	5.94715	5.95225	5.95735	5.96245	5.96755	5.97265	5.97775	5.98285	5.98795
27	5.94210	5.94720	5.95230	5.95740	5.96250	5.96760	5.97270	5.97780	5.98290	5.98800
28	5.94721	5.95231	5.95741	5.96251	5.96761	5.97271	5.97781	5.98291	5.98801	5.99311
29	5.94726	5.95236	5.95746	5.96256	5.96766	5.97276	5.97786	5.98296	5.98806	5.99316
30	5.94731	5.95241	5.95751	5.96261	5.96771	5.97281	5.97791	5.98301	5.98811	5.99321
31	5.94736	5.95246	5.95756	5.96266	5.96776	5.97286	5.97796	5.98306	5.98816	5.99326
32	5.94741	5.95251	5.95761	5.96271	5.96781	5.97291	5.97801	5.98311	5.98821	5.99331
33	5.94746	5.95256	5.95766	5.96276	5.96786	5.97296	5.97806	5.98316	5.98826	5.99336
34	5.94751	5.95261	5.95771	5.96281	5.96791	5.97301	5.97811	5.98321	5.98831	5.99341
35	5.94756	5.95266	5.95776	5.96286	5.96796	5.97306	5.97816	5.98326	5.98836	5.99346
36	5.94761	5.95271	5.95781	5.96291	5.96801	5.97311	5.97821	5.98331	5.98841	5.99351
37	5.94766	5.95276	5.95786	5.96296	5.96806	5.97316	5.97826	5.98336	5.98846	5.99356
38	5.94771	5.95281	5.95791	5.96301	5.96811	5.97321	5.97831	5.98341	5.98851	5.99361
39	5.94776	5.95286	5.95796	5.96306	5.96816	5.97326	5.97836	5.98346	5.98856	5.99366
40	5.94781	5.95291	5.95801	5.96316	5.96826	5.97336	5.97846	5.98356	5.98866	5.99376

## Standard Normal Problem

5.1.1 Suppose that  $Z \sim N(0, 1)$ . Find:

- (a)  $P(Z \leq 1.34)$
- (b)  $P(Z \geq -0.22)$
- (c)  $P(-2.19 \leq Z \leq 0.43)$
- (d)  $P(0.09 \leq Z \leq 1.76)$
- (e)  $P(|Z| \leq 0.38)$
- (f) The value of  $x$  for which  $P(Z \leq x) = 0.55$
- (g) The value of  $x$  for which  $P(Z \geq x) = 0.72$
- (h) The value of  $x$  for which  $P(|Z| \leq x) = 0.31$

Figure 3: image



$$P(Z < 1.34) = \Phi(1.34)$$

$$= .909877$$



$$P(Z > -0.22) = 1 - F(-0.22)$$

$$= 1 - \Phi(-0.22)$$

$$= 1 - .412938$$

$$= .587064$$



## Standard Normal Practice Problems

## Standardizing (the z-transform)

- Suppose  $X$  is a normal random variable with mean  $\mu$  and variance  $\sigma^2$

$$P(X \leq x) = P\left(\frac{X - \mu}{\sigma} \leq \frac{x - \mu}{\sigma}\right) = P(Z \leq z)$$

- The z-value is  $z = (x - \mu)/\sigma$
- Result allows the standard normal tables to be used to calculate probabilities for any normal distribution

## Normal Probability Problem

5.1.3 Suppose that  $X \sim N(10, 2)$ . Find:

- (a)  $P(X \leq 10.34)$
- (b)  $P(X \geq 11.98)$
- (c)  $P(7.67 \leq X \leq 9.90)$
- (d)  $P(10.88 \leq X \leq 13.22)$
- (e)  $P(|X - 10| \leq 3)$
- (f) The value of  $x$  for which  $P(X \leq x) = 0.81$
- (g) The value of  $x$  for which  $P(X \geq x) = 0.04$
- (h) The value of  $x$  for which  $P(|X - 10| \geq x) = 0.63$

Figure 4: image

## Normal Practice Problems

### Normal Approximation to the Binomial Distribution

- If  $X$  is a binomial random variable,

$$Z = \frac{X - np}{\sqrt{np(1-p)}}$$

is approximately a standard normal random variable. Consequently, probabilities computed from  $Z$  can be used to approximate probabilities for  $X$

- Usually holds when

$$np > 5 \quad \text{and} \quad n(1-p) > 5$$

### Problem

4. A process yields 10% defective items. If 100 items are randomly selected from the process, what is the probability that the number of defectives
- (a) exceeds 13?
  - (b) is less than 8?

Source: Walpole, Myers, Myers & Ye

Figure 5: image

- How good are the approximations?

### Continuity Correction Factor

- Is a method to improve the accuracy of the normal approximation to the binomial
- Examine Figure 6.22 from Walpole, Myers, Myers & Ye. Note that each rectangle is centered at  $x$  and extends from  $x - 0.5$  to  $x + 0.5$
- This table should help formulate problems

Binomial Probability	with Correction Factor	Normal Approximation
$P(X \geq x)$	$P(X \geq x - 0.5)$	$P\left(Z > \frac{x-0.5-np}{\sqrt{np(1-p)}}\right)$
$P(X \leq x)$	$P(X \leq x + 0.5)$	$P\left(Z < \frac{x+0.5-np}{\sqrt{np(1-p)}}\right)$
$P(X = x)$	$P(x - 0.5 \leq X \leq x + 0.5)$	$P\left(\frac{x-0.5-np}{\sqrt{np(1-p)}} < Z < \frac{x+0.5-np}{\sqrt{np(1-p)}}\right)$

## Normal Approximation - Figure




Figure 6.22 Normal approximation of  $b(x; 15, 0.4)$ .  
Source: Walpole, Myers, Myers & Ye

Figure 6: image



### Rework Problem using Continuity Correction Factor

- Are the approximations improved?

### Normal Approximation to the Poisson Distribution

- If  $X$  is a Poisson random variable with  $E(X) = \lambda$  and  $V(X) = \lambda$ ,

$$Z = \frac{X - \lambda}{\sqrt{\lambda}}$$

is approximately a standard normal random variable.

## Exponential Distribution

- The exponential distribution is widely used in the area of reliability and life-test data.
- Ostle, et. al. (1996) list the following applications of the exponential distribution
  - the number of feet between two consecutive erroneous records on a computer tape,
  - the lifetime of a component of a particular device,
  - the length of a life of a radioactive material and
  - the time to the next customer service call at a service desk

### Exponential Distribution

- The PDF for an exponential distribution with parameter  $\lambda > 0$  is

$$f(x) = \lambda e^{-\lambda x}, \quad \text{for } 0 \leq x < \infty$$

- The mean of  $X$  is

$$\mu = E(X) = \frac{1}{\lambda}$$

- The variance of  $X$  is

$$\sigma^2 = V(X) = \frac{1}{\lambda^2}$$

Note that other authors define  $f(x) = \frac{1}{\theta} e^{-x/\theta}$ . Either definition is acceptable. However one must be aware of which definition is being used.

### The Exponential CDF

The CDF for the exponential distribution is easy to derive

$$\begin{aligned} F(x) &= P(X \leq x) = \int_{-\infty}^x \lambda e^{-\lambda y} dy \\ &= \int_0^x \lambda e^{-\lambda y} dy \\ &= \left( -e^{-\lambda y} \right) \Big|_{y=0}^x \\ &= -e^{-\lambda x} - (-e^0) \\ &= -e^{-\lambda x} + 1 \\ &= 1 - e^{-\lambda x} \end{aligned}$$

## Problem 4-79

4-79. The time to failure (in hours) of fans in a personal computer can be modeled by an exponential distribution with  $\lambda = 0.0003$ .

- (a) What proportion of the fans will last at least 10,000 hours?
- (b) What proportion of the fans will last at most 7000 hours?

Figure 7: image

### Lack of Memory Property

- The mathematical definition is

$$P(X < t_1 + t_2 | X > t_1) = P(X < t_2)$$

- That is “the probability of a failure time that is less than  $t_1 + t_2$  given the failure time is greater than  $t_1$  is the probability that the item's failure time is less than  $t_2$ ”
- This property is unique to the exponential distribution
- Often used to model the reliability of electronic components.

## Problem 4–80

4-80. The time between the arrival of electronic messages at your computer is exponentially distributed with a mean of two hours.

- (a) What is the probability that you do not receive a message during a two-hour period?
- (b) If you have not had a message in the last four hours, what is the probability that you do not receive a message in the next two hours?
- (c) What is the expected time between your fifth and sixth messages?

Figure 8: image



### Relationship to the Poisson Distribution

- Let  $Y$  be a Poisson random variable with parameter  $\lambda$ . Note:  $Y$  represents the number of occurrences per unit
- Let  $X$  be a random variable that records the time between occurrences for the same process as  $Y$
- $X$  has an exponential distribution with parameter  $\lambda$

## Lognormal Distribution

- Let  $W$  have a normal distribution with mean  $\theta$  and variance  $\omega^2$ ; then  $X = \exp(W)$  is a **lognormal random variable** with pdf

$$f(x) = \frac{1}{x\omega\sqrt{2\pi}} \exp\left[-\frac{(\ln(x) - \theta)^2}{2\omega^2}\right] \quad 0 < x < \infty$$

- The mean of  $X$  is

$$E(X) = e^{\theta + \omega^2/2}$$

- The variance of  $X$  is

$$V(X) = e^{2\theta + \omega^2} (e^{\omega^2} - 1)$$

### Example Problem

3-47. Suppose that  $X$  has a lognormal distribution with parameters  $\theta = 5$  and  $\omega^2 = 9$ . Determine the following:

- (a)  $P(X < 13,300)$
- (b) The value for  $x$  such that  $P(X \leq x) = 0.95$
- (c) The mean and variance of  $X$

Figure 9: image

## Gamma Distribution

- The random variable  $X$  with pdf

$$f(x) = \frac{\lambda^r x^{r-1} e^{-\lambda x}}{\Gamma(r)}, \text{ for } x > 0$$

is a **gamma random variable** with parameters  $\lambda > 0$  and  $r > 0$ .

- The gamma function is

$$\Gamma(r) = \int_0^{\infty} x^{r-1} e^{-x} dx \text{ for } r > 0$$

with special properties:

- $\Gamma(r)$  is finite
- $\Gamma(r) = (r-1)\Gamma(r-1)$
- For any positive integer  $r$ ,  $\Gamma(r) = (r-1)!$

### Gamma Distribution

- The mean and variance are

$$\mu = E(X) = r/\lambda \text{ and } \sigma^2 = V(X) = r/\lambda^2$$

- We will not work any probability problems using the gamma distribution

## Gamma Tables

APPENDIX A

Gamma Percent			
$\alpha$	$T(\alpha)$	$\alpha$	$T(\alpha)$
0.0100	99.4327	0.5100	1.7384
0.0200	99.4423	0.5200	1.7598
0.0300	99.4520	0.5300	1.7814
0.0400	99.4618	0.5400	1.8031
0.0500	99.4716	0.5500	1.8250
0.0600	99.4815	0.5600	1.8470
0.0700	99.4914	0.5700	1.8691
0.0800	99.5013	0.5800	1.8913
0.0900	99.5113	0.5900	1.9136
0.1000	99.5213	0.6000	1.9360
0.1100	99.5313	0.6100	1.9585
0.1200	99.5413	0.6200	1.9811
0.1300	99.5513	0.6300	2.0038
0.1400	99.5613	0.6400	2.0265
0.1500	99.5713	0.6500	2.0493
0.1600	99.5813	0.6600	2.0721
0.1700	99.5913	0.6700	2.0950
0.1800	99.6013	0.6800	2.1179
0.1900	99.6113	0.6900	2.1409
0.2000	99.6213	0.7000	2.1639
0.2100	99.6313	0.7100	2.1869
0.2200	99.6413	0.7200	2.2099
0.2300	99.6513	0.7300	2.2329
0.2400	99.6613	0.7400	2.2559
0.2500	99.6713	0.7500	2.2789
0.2600	99.6813	0.7600	2.3019
0.2700	99.6913	0.7700	2.3249
0.2800	99.7013	0.7800	2.3479
0.2900	99.7113	0.7900	2.3709
0.3000	99.7213	0.8000	2.3939
0.3100	99.7313	0.8100	2.4169
0.3200	99.7413	0.8200	2.4399
0.3300	99.7513	0.8300	2.4629
0.3400	99.7613	0.8400	2.4859
0.3500	99.7713	0.8500	2.5089
0.3600	99.7813	0.8600	2.5319
0.3700	99.7913	0.8700	2.5549
0.3800	99.8013	0.8800	2.5779
0.3900	99.8113	0.8900	2.6009
0.4000	99.8213	0.9000	2.6239
0.4100	99.8313	0.9100	2.6469
0.4200	99.8413	0.9200	2.6699
0.4300	99.8513	0.9300	2.6929
0.4400	99.8613	0.9400	2.7159
0.4500	99.8713	0.9500	2.7389
0.4600	99.8813	0.9600	2.7619
0.4700	99.8913	0.9700	2.7849
0.4800	99.9013	0.9800	2.8079
0.4900	99.9113	0.9900	2.8309
0.5000	99.9213	1.0000	2.8539

## Weibull Distribution

- The random variable  $X$  with pdf

$$f(x) = \frac{\beta}{\delta} \left(\frac{x}{\delta}\right)^{\beta-1} \exp\left[-\left(\frac{x}{\delta}\right)^{\beta}\right], \quad \text{for } x > 0$$

is a **Weibull random variable** with scale parameter  $\delta > 0$  and shape parameter  $\beta > 0$

- The CDF for the Weibull distribution is

$$F(x) = 1 - \exp\left[-\left(\frac{x}{\delta}\right)^{\beta}\right]$$

- The mean of the Weibull distribution is

$$\mu = E(X) = \delta \Gamma\left(1 + \frac{1}{\beta}\right)$$

## Weibull Problem

45. Suppose that fracture strength (MPa) of silicon nitride braze joints under certain conditions has a Weibull distribution with  $\beta = 5$  and  $\sigma = 125$  (suggested by data in the article "Heat-Resistant Active Brazing of Silicon Nitride: Mechanical Evaluation of Braze Joints," (*Welding J.*, August 1997: 300s–304s).
- What proportion of such joints have a fracture strength of at most 100? Between 100 and 150?
  - What strength value separates the weakest 50% of all joints from the strongest 50%?
  - What strength value characterizes the weakest 5% of all joints?

Figure 11: image



## Weibull Practice Problems


## CDF of uniform

Wednesday, February 26, 2025

8:59 AM

$$f(x) = \frac{1}{b-a} \text{ for } a < x < b$$

$$F(x) = \int_{-\infty}^x f(x) dx = \begin{cases} 0 & \text{for } x < a \\ \int_a^x \frac{1}{b-a} dy & \text{for } a < x < b \\ 1 & \text{for } x > b \end{cases}$$

$$\int_a^x \frac{1}{b-a} dy = \left( \frac{1}{b-a} y \right) \Big|_{y=a}^x \rightarrow \frac{x-a}{b-a}$$


$$a = 49.75$$
$$b = 50.25$$


$$P(X > 50) = 1 - F(50)$$
$$= 1 - \frac{50 - 49.75}{50.25 - 49.75} = .5$$

## Part f

Monday, March 3, 2025 8:31 AM

Find  $x$  such that  $P(Z \leq x) = 0.55$   
S.T.

find value in tables that has  
area closest to 0.5 and  
find value of  $z \rightarrow x = 0.3$




## Part g

Monday, March 3, 2025

8:35 AM

Find  $x$  s.t.  $P(Z > x) = .72$

$$x = -.58$$



# Standard normal pp

Monday, March 3, 2025 8:39 AM

## QUESTION 1

Let  $Z$  be a standard normal random variable, find  $P(Z > 2.65)$ .

- ☐ 0.988088
- ☐ The correct answer is not provided.
- ☐ 0.011912
- ☐ 0.137439
- ☐ 0.995975
- ☐ 0.975374
- ☒ 0.004025



$$\begin{aligned} P(Z > 2.65) &= 1 - \Phi(2.65) \\ &= 1 - 0.995975 \\ &= \underline{0.004025} \end{aligned}$$

Screen clipping taken: 3/3/2025 8:40 AM


# Standard normal pp

Monday, March 3, 2025 8:43 AM

## QUESTION 3

Let  $Z$  be a standard normal random variable, find the value  $z$  such that  $P(Z > z) = 0.7825$ .

- ☐ 0.78
- ☐ The correct answer is not provided.
- ☐ 3.12
- ☐ 1.07
- ☒ -0.78
- ☐ -1.07



Screen clipping taken: 3/3/2025 8:43 AM

# Standard normal pp

Monday, March 3, 2025


8:46 AM

## QUESTION 5

Let  $Z$  be a standard normal random variable, find the value  $z$  such that  $P(Z < z) = 0.1986$ .


- ☐ 1.29
- ☐ The correct answer is not provided.
- ☐ 0.85
- ☐ 3.1
- ☐ -1.29
- ☒ -0.85

Screen clipping taken: 3/3/2025 8:46 AM



# Standard normal pp

Monday, March 3, 2025 8:50 AM



Let  $Z$  be a standard normal random variable, find  $P(1.22 < Z < 1.28)$ .

☐ 0.010959

☐ -0.013696

☒ The correct answer is not provided.

☐ 0.100273

☐ 0.888768

☐ 0.989041

$$\begin{aligned} P(1.22 < Z < 1.28) &= \Phi(1.28) - \Phi(1.22) \\ &= .899727 - .888767 \\ &= \cancel{.01096} \\ &= \underline{\underline{.01096}} \end{aligned}$$

Screen clipping taken: 3/3/2025 8:50 AM




# Standard normal pp

Monday, March 3, 2025 8:56 AM

## QUESTION 9

Let  $Z$  be a standard normal random variable, find  $P(Z < 3.07)$ .

- ☐ 0.003584
- ☐ 0.996416
- ☐ 0.00107
- ☐ The correct answer is not provided.
- ☐ 0.99893
- ☐ 0.878258
- ☐ 0.839243



$$P(Z < 3.07) = \Phi(3.07) = .998930$$

Screen clipping taken: 3/3/2025 8:56 AM

Attended

line 1 E)

line 2 A) March 24 , B) March 26  
Monday Wednesday